

LAKE: KIDNEY P
 TOWN: T03 R10 WELS
 COUNTY: PISCATAQUIS

MIDAS: 716
 TRUE BASIN: 1
 SAMPLE STATION: 1

WHOLE LAKE INFORMATION



MAX. DEPTH: 10 m. (33 ft.)
 MEAN DEPTH: 3 m. (10 ft.)
 DELORME ATLAS #: 50
 USGS QUAD: DOUBLETOP MOUNTAIN
 IFW REGION F: Penobscot (Enfield)
 IFW FISH. MANAGMENT: Coldwater

TRUE BASIN CHARACTERISTICS

SURFACE AREA: 37.0 ha. (91.4 a.)
 FLUSHING RATE: 2.01 flushes/yr.
 VOLUME: 921720.0 cu. m. (748 ac.-ft.)
 DIRECT DRAINAGE AREA: 1.43 sq. km. (0.55 sq. mi.)

PLEASE NOTE THE FOLLOWING: The SAMPLE STATION # refers to the location sampled. The term TRUE BASIN is used to define areas within a lake that are separated by shallow reefs or shoals and therefore function as separate lakes. There are approximately 50 lakes in the state that have more than 1 True Basin. True Basin Characteristics are now being included in the first section of these reports to enable users of the Phosphorous Loading Methodology to better evaluate the data. If there is no data for a particular True Basin, True Basin Characteristics must be obtained from the DEP. KIDNEY P has 1 True Basin(s).

SECCHI DISK TRANSPARENCY GRAPHS:



Note: 2009 graphs may indicate multiple readings taken on a given day.

SUMMARY OF CHEMICAL AND TROPHIC STATE PARAMETERS:

[* indicates that Secchi disk was visible at bottom of lake (or one reading used in calculation was visible)].

YEAR	MEAN COLOR (SPU)	MEAN pH	MEAN ALK (mg/l)	MEAN COND. (uS/cm)	TOTAL PHOS. MEANS (ppb)				SECCHI DISK (m.)				CHLOROPHYLL A(ppb)			TROPIC STATE INDICE			
					EPI	SURF	BOT.	PRO.	MIN.	MEAN	MAX.	N	MIN.	MEAN	MAX.	C	G	SEC	CH
1981	25	6.80	8.0	26	8	-	-	13	4.9	5.2	5.8	2	2.7	2.7	2.7	-	-	-	-
1982	-	-	-	-	-	-	-	-	4.9	5.2	5.3	3	-	-	-	-	-	-	-
1983	-	-	-	-	-	-	-	-	5.0	5.5	5.9	3	-	-	-	-	-	-	-
1984	30	7.00	6.0	-	9	-	-	41	3.5	5.4	7.3	4	2.6	2.6	2.6	-	-	-	-
1992	-	-	-	-	-	-	-	-	4.6	4.6	4.6	1	-	-	-	-	-	-	-
1994	-	-	-	-	-	-	-	-	4.5	4.5	4.5	1	-	-	-	-	-	-	-
1995	-	-	-	-	-	-	-	-	5.5	5.5	5.5	1	-	-	-	-	-	-	-
1996	16	-	6.0	25	7	-	-	16	4.1	4.3	4.5	1	2.8	2.8	2.8	-	-	-	-
-	-	-	-	-	-	-	-	-	4.5	4.5	4.5	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	5.5	5.5	5.5	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	4.0	4.0	4.0	1	-	-	-	-	-	-	-
2003	-	-	-	-	-	-	-	-	4.5	4.5	4.5	1	-	-	-	-	-	-	-
2008	-	-	-	-	-	-	-	-	3.1	3.1	3.1	1	-	-	-	-	-	-	-
2009	-	-	-	-	-	-	-	-	4.0	4.0	4.0	1	-	-	-	-	-	-	-
SUMMARY:	24	6.89	6.7	26	8	-	-	23	3.1	4.7	7.3	14	2.6	2.7	2.8	-	-	-	-

LAKE: KIDNEY P
TOWN: T03 R10 WELS
COUNTY: PISCATAQUIS

MIDAS: 716
*TRUE BASIN: 1
*SAMPLE STATION: 1

LATE SUMMER TEMPERATURE / DISSOLVED OXYGEN PROFILES:

DEPTH	SAMPLE DATE							
	09/01/81		08/27/84		08/21/96		08/31/96	
m	°C	ppm	°C	ppm	°C	ppm	°C	ppm
0.0	19.0	8.7	-	-	21.2	8.2	19.5	0.0
1.0	19.5	8.5	22.0	9.5	21.2	8.3	-	-
2.0	19.5	8.5	20.5	9.7	20.4	8.2	-	-
3.0	19.2	8.3	20.0	9.5	19.7	7.4	-	-
4.0	18.5	7.4	18.8	8.0	17.1	1.8	-	-
5.0	17.5	3.5	15.5	1.9	14.1	0.2	-	-
6.0	15.0	0.5	13.0	0.1	11.0	0.1	-	-
7.0	12.0	0.5	10.9	0.1	9.9	0.1	-	-
8.0	10.0	0.5	9.9	0.1	9.0	0.1	-	-
9.0	10.0	0.5	9.8	0.1	8.8	0.1	-	-
10.0	9.0	0.4	9.8	0.1	8.8	0.1	-	-
11.0	9.0	0.3	-	-	-	-	-	-
12.0	9.0	0.3	-	-	-	-	-	-

WATER QUALITY SUMMARY

KIDNEY POND, T03 R10 Wels

Midas: 716, Sample Station # 1

The Maine Department of Environmental Protection (ME-DEP) and the Volunteer Lake Monitoring Program (VLMP) have collaborated in the collection of lake data to evaluate present water quality, track algae blooms, and determine water quality trends. This dataset does not include bacteria, mercury, or nutrients other than phosphorus.

Water quality monitoring data for Kidney Pond has been collected since 1981. During this period, 3 years of basic chemical information was collected, in addition to Secchi Disk Transparencies (SDT). In summary, the water quality of Kidney Pond is considered to be slightly above average, based on measures of SDT, total phosphorus (TP), and Chlorophyll-a (Chla). The potential for nuisance alga blooms on Kidney Pond is low.

Water Quality Measures: Kidney Pond is a non-colored lake (average color 24 SPU) with an average SDT of 4.9m (16.7ft). The range of water column TP for Kidney Pond is 7-9 parts per billion (ppb) with an average of 8 ppb (State average is 12 ppb), while Chla ranges from 2.6-2.8 ppb with an average of 2.7 ppb (State average is 4.8 ppb). Recent dissolved oxygen (DO) profiles show low to moderate DO depletion in deep areas of the lake. The potential for TP to leave the bottom sediments and become available to algae in the water column (internal loading) is low to moderate. Oxygen levels below 5 parts per million stress certain cold water fish, and a persistent loss of oxygen may eliminate or reduce habitat for sensitive cold water species.

Kidney Pond lies within the boundaries of Baxter State Park.

See the ME-DEP *Explanation of Lake Water Quality Monitoring Report* for measured variable explanations. Additional lake information can be obtained by contacting the Maine DEP at 207-287-3901 or the VLMP at 207-783-7733, or on the Internet at <http://www.pearl.maine.edu> and/or <http://www.maine.gov/dep/blwq/lake.htm>.

Filename: kidn716, Revised: 2/02, By: mg